

Medvedev degrees of generalized r.e. separating classes

Douglas Cenzer and Peter G. Hinman

Important examples of Π_1^0 classes of functions $f \in {}^\omega\omega$ are the classes of *sets* (elements of ${}^\omega 2$) which separate a given pair of disjoint r.e. sets: $S_2(A_0, A_1) := \{f \in {}^\omega 2 : (\forall i < 2)(\forall x \in A_i)f(x) \neq i\}$. A wider class consists of the classes of functions $f \in {}^\omega k$ which in a generalized sense separate a k -tuple of r.e. sets (not necessarily pairwise disjoint) for each $k \in \omega$: $S_k(A_0, \dots, A_{k-1}) := \{f \in {}^\omega k : (\forall i < k)(\forall x \in A_i)f(x) \neq i\}$. We study the structure of the Medvedev degrees of such classes and show that the set of degrees realized depends strongly on both k and the extent to which the r.e. sets intersect. Let \mathcal{S}_k^m denote the Medvedev degrees of those $S_k(A_0, \dots, A_{k-1})$ such that no $m+1$ sets among A_0, \dots, A_{k-1} have a nonempty intersection. It is shown that each \mathcal{S}_k^m is an upper semi-lattice but not a lattice. The degree of the set of k -ary diagonally nonrecursive functions DNR_k is the greatest element of \mathcal{S}_k^1 . If $2 \leq l < k$, then $\mathbf{0}_M$ is the only degree in \mathcal{S}_l^1 which is below a member of \mathcal{S}_k^1 . Each \mathcal{S}_k^m is densely ordered and has the splitting property and the same holds for the lattice \mathcal{L}_k^m it generates. The elements of \mathcal{S}_k^m are exactly the joins of elements of \mathcal{S}_i^1 for $\lceil \frac{k}{m} \rceil \leq i \leq k$.